

Possible compensatory role among chloroplast proteases under excess-light stress condition

Yusuke Kato and Wataru Sakamoto*

Institute of Plant Science and Resources (IPSR); Okayama University; Kurashiki, Okayama, Japan

Keywords: D1 protein degradation, Deg, FtsH, photoinhibition, photosynthesis, photosystem II repair cycle, protease, chloroplast

Abbreviations: Col, Columbia; LHCII, light-harvesting complex of photosystem II; PSII, photosystem II; Rubisco, ribulose-1,5-bisphosphate carboxylase/oxygenase; *var2*, *yellow variegated2*

The reaction center protein D1 of photosystem II (PSII), known as a primary target of photodamage, is repaired efficiently by the *PSII repair cycle*, to cope with constant photooxidative damage. Recent studies of *Arabidopsis* show that the endo-type Deg protease and the exo-type FtsH proteases cooperatively degrade D1 in the PSII repair in vivo. It is particularly interesting that we observed upregulation of Clp and SppA proteases when FtsH was limited in the mutant lacking FtsH2. To examine how the complementary functions of chloroplastic proteases are commonly regulated, we undertook a high-light stress on wild-type *Arabidopsis* leaves. The result that wild type leaves also showed increased levels of these proteases upon exposure to excessively strong illumination not only revealed the importance of FtsH and Deg in the PSII repair, but also implied cooperation among chloroplastic proteases under chronic stress conditions.

Light energy constantly damages photosynthetic apparatus, and frequently engenders an inhibitory effect on photosynthesis, known as photoinhibition.¹ Photosynthetic organisms have therefore developed various systems that minimize damage to photosynthetic apparatus and which avoid the accumulation of damaged proteins.^{2,3} Because a primary target of the damage is the photosystem II (PSII) reaction center protein D1, quality control of PSII (i.e., rapid and specific replacement of photodamaged D1 with de-novo synthesized D1 in the PSII repair cycle) is an important mechanism.^{4,5} Numerous studies of the PSII repair cycle have elucidated rapid D1 turnover. In addition, recent studies have revealed that proteases of two types, FtsH and Deg, participate in this process in chloroplasts.^{6,7}

FtsH and Deg have been extensively characterized in *Arabidopsis*. FtsH is a zinc metalloprotease that processively degrades membrane proteins.⁸ In chloroplasts, FtsH forms a hetero-hexameric complex.⁹⁻¹¹ The *Arabidopsis* mutants lacking a major isoform FtsH2 or FtsH5, which constitutes two representative isoforms in the heterocomplex, show leaf-variegated phenotypes (termed *yellow variegated2*, *var2* and *var1*, respectively) and accumulate high levels of reactive oxygen species.¹²⁻¹⁴ Significant impairment of the rate of D1 degradation in the mutant lacking FtsH provided compelling evidence that FtsH is critically important for D1 degradation in the PSII repair.¹⁵⁻¹⁷ As for Deg, five isoforms are identified in *Arabidopsis* chloroplasts: Deg1, Deg5 and Deg8 are localized on the luminal side, whereas Deg2 and Deg7 are on the stromal side.¹⁸⁻²¹ Deg proteases are

regarded as cleaving D1 at surface-exposed loops between transmembrane helices under photoinhibitory light conditions.

Our recent study demonstrated that the fragmented D1 polypeptides, which were detectable only in *var2* background under excessively strong illumination (2,500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$), resulted from Deg protease activities.²² These results thus provided in vivo evidence that FtsH and Deg degrade photodamaged D1 cooperatively under photoinhibitory conditions, as shown by in vitro studies. We proposed a model by which, in chloroplasts, fundamental D1 degradation is conducted by FtsH at all light intensities and that Deg proteases assist effective D1 degradation by producing additional recognition termini that are accessible to FtsH under high light.

In addition to cooperation between FtsH and Deg in PSII repair, one noteworthy finding was upregulation of several other chloroplastic proteases in *var2*.²² Clp is a stromal multi-subunit processive protease that is structurally and functionally related to FtsH. Our recent study demonstrated that Clp levels increased in *var2*. Similarly, a thylakoid membrane-bound protease, SppA, was shown to be upregulated in *var2*. It is particularly interesting that a portion of Clp was recruited from stroma to the thylakoid membrane. Given the functional similarity between Clp and FtsH,²³ it is plausible that Clp can be partially substituted for FtsH, suggesting compensation among chloroplastic proteases.

To examine how the complementary functions of chloroplastic proteases are commonly regulated, here we used wild type

*Correspondence to: Wataru Sakamoto; Email: saka@rib.okayama-u.ac.jp

Submitted: 11/09/2012; Revised: 12/10/2012; Accepted: 12/10/2012

<http://dx.doi.org/10.4161/psb.23198>

Citation: Kato Y, Sakamoto W. Possible compensatory role among chloroplast proteases under excess-light stress condition. Plant Signal Behav 2013; 8: e23198.




Figure 1. Accumulation and localization of chloroplast proteases under growth light condition and after excessively strong illumination. Chloroplasts were purified using a Percoll step gradient from mature leaves of Col (ca. 6-week-old plants) that were illuminated in growth light (GL) ($180 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) and excessively strong illumination conditions (HL) ($2,500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) for 6 h. Intact chloroplasts were fractionated into stroma and membrane fractions, according to the protocol as reported previously.²² A representative immunoblot using specific antibodies (FtsH2/8, ClpP6, ClpC1/2, SppA, Deg2, Deg8, D1 and Rubisco large subunit) and the bands corresponding to CBB-stained LHCII are shown. D1 and Rubisco large subunit were used, respectively, as markers of membranes and stroma. Samples were equally loaded based on chlorophyll content. These experiments were performed three times with similar results.

Columbia (Col) plants instead of *var2*, and observed accumulation of Clp and SppA under excessively strong illumination. Immunoblotting using antibodies against these proteases was performed along with protein fractions from total chloroplasts, stroma and thylakoid membranes. Except for the use of Col and a different light intensity, other experimental condition was identical to those reported in our previous work.²² First, we compared levels of these proteases after short-time excess light exposure ($2,500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ for 1 h). A significant change in the protease level, however, was never observed. This result made us postulate that constitutive accumulation of damaged proteins in thylakoid membranes, most likely as represented by

the situation in *var2*, triggers upregulation of Clp and SppA. To induce constitutive accumulation of damaged proteins in Col, we next examined a chronic high-light stress by prolonged exposure at excess light ($2,500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ for 6 h). As a consequence of this exposure, maximum quantum yield of PSII (Fv/Fm) in Col leaves was decreased to approximately 40% of the initial value before light illumination (Fv/Fm before and after light illumination was 0.83 ± 0.01 and 0.36 ± 0.01 , respectively). We found that the prolonged exposure for 6 h caused increased accumulation of Clp and SppA and recruited stromal Clp to thylakoid membranes (Fig. 1). These results are consistent with our observation in *var2*, indicating that the compensatory regulation of chloroplast proteases occurs in the wild type.²² In contrast, we found no significant difference in Deg2 and Deg8 levels in thylakoid membranes after the same light exposure. We observed considerable decrease in D1 protein after the same light exposure, suggesting the excess damage on photosynthetic apparatuses. Whereas, no significant changes of FtsH, Rubisco large subunit and LHC proteins were detected.

It is noteworthy that SppA increased dramatically after treatment, which is consistent with a previous observation that SppA is upregulated by strong illumination.²⁴ Overall, these results confirm that chronic damage to thylakoid membranes causes changes in the accumulation level of Clp and SppA proteases and the membrane localization of Clp, which implicates the compensation of the chloroplastic proteases in degrading D1 and possibly other proteins in the thylakoid membrane. Although further investigation is necessary, long-term acclimation to light stress and chronic damage to thylakoid membranes might be likely triggers of cooperation among chloroplastic proteases. Consequently, our recent work not only explains cooperative degradation of D1 by FtsH and Deg in the PSII repair; it also elucidates the compensatory control of chloroplast proteases related to high-light acclimation.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

This work was supported by the Japan Society for the Promotion of Science [Grant-in-Aid for Young Scientists (No. 22770042 to Y.K.)], Japan Science and Technology Agency [Core Research for Evolutional Science and Technology (to W.S.)] and the Oohara Foundation (to W.S.).

References

1. Aro EM, Virgin I, Andersson B. Photoinhibition of Photosystem II. Inactivation, protein damage and turnover. *Biochim Biophys Acta* 1993; 1143:113-34; PMID:8318516; [http://dx.doi.org/10.1016/0005-2728\(93\)90134-2](http://dx.doi.org/10.1016/0005-2728(93)90134-2).
2. Murchie EH, Niyogi KK. Manipulation of photoprotection to improve plant photosynthesis. *Plant Physiol* 2011; 155:86-92; PMID:21084435; <http://dx.doi.org/10.1104/pp.110.168831>.
3. Takahashi S, Badger MR. Photoprotection in plants: a new light on photosystem II damage. *Trends Plant Sci* 2011; 16:53-60; PMID:21050798; <http://dx.doi.org/10.1016/j.tplants.2010.10.001>.
4. Edelman M, Mattoo AK. D1-protein dynamics in photosystem II: the lingering enigma. *Photosynth Res* 2008; 98:609-20; PMID:18709440; <http://dx.doi.org/10.1007/s11120-008-9342-x>.
5. Nixon PJ, Michoux F, Yu J, Boehm M, Komenda J. Recent advances in understanding the assembly and repair of photosystem II. *Ann Bot* 2010; 106:1-16; PMID:20338950; <http://dx.doi.org/10.1093/aob/mcq059>.
6. Kato Y, Sakamoto W. Protein quality control in chloroplasts: a current model of D1 protein degradation in the photosystem II repair cycle. *J Biochem* 2009; 146:463-9; PMID:19451147; <http://dx.doi.org/10.1093/jb/mvp073>.
7. Komenda J, Sobotka R, Nixon PJ. Assembling and maintaining the Photosystem II complex in chloroplasts and cyanobacteria. *Curr Opin Plant Biol* 2012; 15:245-51; PMID:22386092; <http://dx.doi.org/10.1016/j.pbi.2012.01.017>.
8. Ito K, Akiyama Y. Cellular functions, mechanism of action, and regulation of FtsH protease. *Annu Rev Microbiol* 2005; 59:211-31; PMID:15910274; <http://dx.doi.org/10.1146/annurev.micro.59.030804.121316>.
9. Lindahl M, Tabak S, Cseke L, Pichersky E, Andersson B, Adam Z. Identification, characterization, and molecular cloning of a homologue of the bacterial FtsH protease in chloroplasts of higher plants. *J Biol Chem* 1996; 271:29329-34; PMID:8910594; <http://dx.doi.org/10.1074/jbc.271.46.29329>.

10. Sakamoto W, Zaltsman A, Adam Z, Takahashi Y. Coordinated regulation and complex formation of *yellow variegated1* and *yellow variegated2*, chloroplastic FtsH metalloproteases involved in the repair cycle of photosystem II in *Arabidopsis* thylakoid membranes. *Plant Cell* 2003; 15:2843-55; PMID:14630971; <http://dx.doi.org/10.1105/tpc.017319>.
11. Yu F, Park S, Rodermeier SR. The *Arabidopsis* FtsH metalloprotease gene family: interchangeability of subunits in chloroplast oligomeric complexes. *Plant J* 2004; 37:864-76; PMID:14996218; <http://dx.doi.org/10.1111/j.1365-3113X.2003.02014.x>.
12. Chen M, Choi Y, Voytas DF, Rodermeier S. Mutations in the *Arabidopsis* *VAR2* locus cause leaf variegation due to the loss of a chloroplast FtsH protease. *Plant J* 2000; 22:303-13; PMID:10849347; <http://dx.doi.org/10.1046/j.1365-3113x.2000.00738.x>.
13. Takeuchi K, Sodmergen, Murata M, Motoyoshi F, Sakamoto W. The *YELLOW VARIEGATED (VAR2)* locus encodes a homologue of FtsH, an ATP-dependent protease in *Arabidopsis*. *Plant Cell Physiol* 2000; 41:1334-46; PMID:11134419; <http://dx.doi.org/10.1093/pcp/pcd067>.
14. Sakamoto W, Tamura T, Hanba-Tomita Y, Murata M, Sodmergen. The *VARI* locus of *Arabidopsis* encodes a chloroplastic FtsH and is responsible for leaf variegation in the mutant alleles. *Genes Cells* 2002; 7:769-80; PMID:12167156; <http://dx.doi.org/10.1046/j.1365-2443.2002.00558.x>.
15. Kato Y, Sakamoto W. Protein quality control in chloroplasts: a current model of D1 protein degradation in the photosystem II repair cycle. *J Biochem* 2009; 146:463-9; PMID:19451147; <http://dx.doi.org/10.1093/jb/mvp073>.
16. Bailey S, Thompson E, Nixon PJ, Horton P, Mullineaux CW, Robinson C, et al. A critical role for the Var2 FtsH homologue of *Arabidopsis thaliana* in the photosystem II repair cycle *in vivo*. *J Biol Chem* 2002; 277:2006-11; PMID:11717304; <http://dx.doi.org/10.1074/jbc.M105878200>.
17. Clausen T, Southan C, Ehrmann M. The HtrA family of proteases: implications for protein composition and cell fate. *Mol Cell* 2002; 10:443-55; PMID:12408815; [http://dx.doi.org/10.1016/S1097-2765\(02\)00658-5](http://dx.doi.org/10.1016/S1097-2765(02)00658-5).
18. Haussühl K, Andersson B, Adamska I. A chloroplast DegP2 protease performs the primary cleavage of the photodamaged D1 protein in plant photosystem II. *EMBO J* 2001; 20:713-22; PMID:11179216; <http://dx.doi.org/10.1093/emboj/20.4.713>.
19. Kapri-Pardes E, Naveh L, Adam Z. The thylakoid lumen protease Deg1 is involved in the repair of photosystem II from photoinhibition in *Arabidopsis*. *Plant Cell* 2007; 19:1039-47; PMID:17351117; <http://dx.doi.org/10.1105/tpc.106.046573>.
20. Sun X, Peng L, Guo J, Chi W, Ma J, Lu C, et al. Formation of DEG5 and DEG8 complexes and their involvement in the degradation of photodamaged photosystem II reaction center D1 protein in *Arabidopsis*. *Plant Cell* 2007; 19:1347-61; PMID:17449806; <http://dx.doi.org/10.1105/tpc.106.049510>.
21. Sun X, Fu T, Chen N, Guo J, Ma J, Zou M, et al. The stromal chloroplast Deg7 protease participates in the repair of photosystem II after photoinhibition in *Arabidopsis*. *Plant Physiol* 2010; 152:1263-73; PMID:20089771; <http://dx.doi.org/10.1104/pp.109.150722>.
22. Kato Y, Sun X, Zhang L, Sakamoto W. Cooperative D1 degradation in the photosystem II repair mediated by chloroplastic proteases in *Arabidopsis*. *Plant Physiol* 2012; 159:1428-39; PMID:22698923; <http://dx.doi.org/10.1104/pp.112.199042>.
23. Kato Y, Sakamoto W. New insights into the types and function of proteases in plastids. *Int Rev Cell Mol Biol* 2010; 280:185-218; PMID:20797683; [http://dx.doi.org/10.1016/S1937-6448\(10\)80004-8](http://dx.doi.org/10.1016/S1937-6448(10)80004-8).
24. Wetzel CM, Harmacek LD, Yuan LH, Wopereis JL, Chubb R, Turini P. Loss of chloroplast protease SPPA function alters high light acclimation processes in *Arabidopsis thaliana* L. (Heynh.). *J Exp Bot* 2009; 60:1715-27; PMID:19349419; <http://dx.doi.org/10.1093/jxb/erp051>.